

CLINICAL CASE SERIES

The Impact of Diabetes on the Outcomes of Surgical and Nonsurgical Treatment of Patients in the Spine Patient Outcomes Research Trial

Mitchell K. Freedman, DO,* Alan S. Hilibrand, MD,* Emily A. Blood, PhD,† Wenyan Zhao, MS,† Todd J. Albert, MD,* Alexander R. Vaccaro, MD, PhD,* Christina V. Oleson, MD,‡ Tamara S. Morgan, MA,†‡ and James N. Weinstein, MD†‡

Study Design. A secondary analysis comparing diabetic patients with nondiabetic patients enrolled in the Spine Patient Outcomes Research Trial (SPORT).

Objective. To compare surgical outcomes and complications between diabetic and nondiabetic spine patients.

Summary of Background Data. Patients with diabetes are predisposed to comorbidities that may confound the diagnosis and treatment of patients with spinal disorders.

Methods. Baseline characteristics and outcomes of 199 patients with diabetes were compared with those of the nondiabetic population in a total of 2405 patients enrolled in the Spine Patient Outcomes Research Trial for the diagnoses of intervertebral disc herniation (IDH), spinal stenosis (SpS), and degenerative spondylolisthesis (DS). Primary outcome measures include the 36-Item Short Form Health Survey (SF-36) Health Status questionnaire and the Oswestry Disability Index.

Results. Patients with diabetes were significantly older and had a higher body mass index than nondiabetic patients. Comorbidities, including hypertension, stroke, cardiovascular disease, and joint disease, were significantly more frequent in diabetic patients than in nondiabetic patients. Patients with diabetes and IDH did not make significant gains in pain and function with surgical intervention relative to diabetic patients who underwent nonoperative treatment. Diabetic patients with SpS and DS experienced significantly greater improvements in pain and function with surgical intervention when compared with nonoperative treatment. Among those who had surgery, nondiabetic patients with SpS achieved marginally

significantly greater gains in function than their diabetic counterparts (SF-36 physical function, $P = 0.062$). Among patients who had surgery for DS, diabetic patients did not have as much improvement in pain or function as did the nondiabetic population (SF-36 bodily pain, $P = 0.003$; physical function, $P = 0.002$). Postoperative complications were more prevalent in patients with diabetes than in nondiabetic patients with SpS ($P = 0.002$). There was an increase in postoperative ($P = 0.028$) and intraoperative ($P = 0.029$) blood replacement in DS patients with diabetes.

Conclusion. Diabetic patients with SpS and DS benefited from surgery, though older SpS patients with diabetes have more postoperative complications. IDH patients with diabetes did not benefit from surgical intervention.

Key words: complications, degenerative spondylolisthesis, diabetes mellitus, disability, intervertebral disc herniation, pain, spinal stenosis.

Spine 2011;36:290–307

Diabetes mellitus is a debilitating chronic illness that affects 16 million Americans.¹ Although the disease itself is not necessarily disabling, many of its sequelae, including diabetic neuropathy and microvascular disease, can cause chronic lower extremity pain and lead to significant limitation in overall function.^{1–7} The coexistence of diabetic and lumbar spine disease may cause even greater limitation among the diabetic population when compared with nondiabetic population, prompting more aggressive treatment.

The literature contains conflicting reports regarding the benefits of surgical decompression and fusion among diabetic patients with lumbar spinal stenosis (SpS) and degenerative disc disease.^{8–12} In addition, it has been suggested that patients with diabetes may be predisposed to complications, such as infection, prolonged hospitalization, longer operative time, and higher nonunion rate, after spinal surgery.^{12–16} After surgical intervention, patients with diabetes may also have poor health status and decreased life expectancy compared with nondiabetic patients.^{17,18}

This study compares the baseline characteristics of patients with diabetes to the nondiabetic patients in the Spine Patient Outcomes Research Trial (SPORT). The study also evaluates the impact of the diabetic condition on the clinical outcomes of operative and nonoperative treatment.^{19–23}

From the *Rothman Institute, Thomas Jefferson University, Philadelphia, PA; †the Dartmouth Institute for Health Policy and Clinical Practice and Dartmouth Medical School, Hanover, NH; and ‡Department of Orthopaedics, Dartmouth-Hitchcock Medical Center, Lebanon, NH.

Acknowledgement date: August 19, 2009. Revision date: November 30, 2009. Acceptance date: December 1, 2009.

The manuscript submitted does not contain information about medical device(s)/drug(s).

No funds were received in support of this work. No benefits in any form have been or will be received from a commercial party related directly or indirectly to the subject of this manuscript.

Address correspondence and reprint requests to Mitchell Freedman, DO, Rothman Institute, Thomas Jefferson University, Philadelphia, PA 19107; E-mail: mitchell.freedman@rothmaninstitute.com

DOI: 10.1097/BRS.0b013e3181ef9d8c

MATERIALS AND METHODS

SPORT enrolled patients from 2000 to 2005 at 13 sites in 11 US states. Subjects were enrolled in both observational and randomized cohorts for all diagnoses. Patients chose whether or not they were willing to be randomized. If they were, they were enrolled in the randomized cohort. If they were not, they were enrolled in the observational cohort. Once in the randomized cohort, they were randomized to surgery or nonoperative treatment. Once in the observational cohort, they elected to have surgery or nonoperative treatment. The primary outcome measures were the 36-Item Short Form Health Survey (SF-36) Health Status questionnaire subscores for bodily pain (BP) and physical function (PF),^{24,25} as well as low back pain-associated disability as measured by the Oswestry Disability Index (ODI).²⁶ Baseline characteristics analyzed in this study included age, sex, race, income, work status, disability status, medical comorbidities, mean body mass index (BMI), smoking, and health status, as measured by SF-36. Further details about the SPORT study have been previously published.^{19–23,27,28}

There are a total of 2505 patients enrolled in the observational and randomized cohorts of the SPORT trial, 2406 of whom provided information regarding diabetic status. The cohorts were combined after analyzing each separately and testing for differences in effect sizes between randomized and observational cohorts. Since no differences were seen, the cohorts were combined for as-treated analyses. This was done in several previous SPORT secondary analysis articles,^{29–31} and a detailed statistical rationale for this strategy has been published.³²

Baseline characteristics for all patients for whom follow-up and diabetes status were available were compiled and comparisons made between diabetic patients and nondiabetic patients (Table 1). All 2505 enrolled patients completed a baseline survey. If a patient completed a baseline survey, they might have missing values for those questions they refused to answer. Therefore, the number of patients varied with each baseline characteristic. These numbers, along with the total number of patients included in the analysis, are included in Table 1. No patients were missing age. Only patients who had both baseline and at least one follow-up were included in the analyses. Only two patients were missing BMI; they both were in the intervertebral disc herniation (IDH), nondiabetic group. Further, the diabetic and nondiabetic groups were stratified by diagnosis (IDH, SpS, and degenerative spondylolisthesis [DS]).

STATISTICAL ANALYSIS

Baseline characteristics between patients with diabetes and patients without diabetes were compared by using a χ^2 test for categorical variables and t tests for continuous variables. Outcome analyses were performed, as they were in the primary SPORT articles for the individual diagnoses.^{19–21,27,28} Outcomes were analyzed by using longitudinal mixed-effects models, with a random individual effect to account for the correlation among repeated observations within individuals over time. Adjusting covariates found to predict missing data, treatment received, and outcome were included in the model (further details described previously).^{19–21,23,27} In addition, outcome, center, age, and sex were included in all longitudinal

outcome models. All analyses were as-treated, and treatment is considered a time-varying covariate. Therefore, patients were categorized at each time-point as to whether or not they received surgical treatment; follow-up times were measured from the beginning of treatment, and baseline covariates were updated at the time of surgery. All observations before surgery were considered in the nonoperative estimate, with follow-up time measured from enrollment; all observations after surgery contributed to the surgical estimate, with follow-up time measured from the time of surgery. Rates of repeated surgery at 1, 2, 3, and 4 years were estimated via Kaplan-Meier curves. P values were calculated with the use of the log-rank test. Secondary and binary outcomes were analyzed by using generalized estimating equations, assuming a compound symmetry working correlation structure. Comparisons in outcomes between patients with diabetes and patients without diabetes are made at each time-point with multiple degree-of-freedom Wald tests. Across the 4-year follow-up, overall comparisons of area under the curve were made by using a Wald test. Throughout the text, we have distinguished between these by using the terms “at XX time” *versus* “across 4 years.” Analyses were performed with the SAS PROC MIXED and PROC GENMOD procedures (SAS version 9.2, Windows ZP Pro, Cary, NC). *Statistical significance* is defined as $P < 0.05$ based on a two-sided hypothesis, with no adjustment made for multiple comparisons.

RESULTS

Information about follow-up outcomes and diabetic status was available for 2406 patients who were included in the analysis (Table 1). Please note that we here discuss only those differences that pertain to diabetes; those significant differences that remain have been reported and discussed in previous SPORT articles.

One hundred ninety-nine patients reported being treated for diabetes having been told that they had diabetes by their physician. The mean age of the patients with diabetes (63.9 years) was significantly greater than that of the nondiabetic patients (52.8 years) ($P < 0.001$). A significantly higher percentage of patients with diabetes were nonwhites (23% *vs.* 14%, $P < 0.001$). Patients with diabetes had a significantly higher mean BMI than nondiabetic patients (32.9 *vs.* 28.3, $P < 0.001$). Several medical conditions were significantly more common among diabetic patients, including hypertension (64% *vs.* 27%, $P < 0.001$), stroke (6% *vs.* 1%, $P < 0.001$), cardiac conditions (30% *vs.* 13%, $P < 0.001$), and “joint” problems (55% *vs.* 36%, $P < 0.001$). There was also a significantly higher incidence of lung cancer and stomach problems among patients with diabetes. Vascular problems were more prevalent in patients with diabetes.

Significantly, more nondiabetic patients were working (48%) than were the patients with diabetes (26%), and more patients with diabetes were disabled (16%) than nondiabetic patients (11%). Among the SF-36 subscales, there were significant differences between the groups at baseline with respect to the physical component summary (PCS) score (27.1 among patients with diabetes *vs.* 30.3 for nondiabetic patients,

TABLE 1. Patient Baseline Demographic Characteristics, Comorbidities, and Health Status Measures

Characteristics	All Analyzed Patients (IDH + SpS + DS)			IDH		
	Not Diabetic, n = 2207	Diabetic,* n = 199	P†	Not Diabetic, n = 1145	Diabetic,* n = 40	P†
Age, yr; mean (SD)	52.8 (16.3)	63.9 (11.9)	<0.001	41.5 (11.3)	50.4 (11.8)	<0.001
Female, n (%)	1061 (48)	99 (50)	0.70	488 (43)	17 (42)	0.88
Ethnicity, not Hispanic, n (%)‡	2117 (96)	193 (97)	0.59	1094 (96)	37 (92)	0.60
Race, white, n (%)§	1900 (86)	153 (77)	<0.001	995 (87)	31 (78)	0.14
Education, at least some college; n (%)	1548 (70)	121 (61)	0.008	852 (74)	26 (65)	0.25
Income, <\$50,000; n (%)	740 (34)	51 (26)	0.028	513 (45)	19 (48)	0.86
Marital Status, married; n (%)	1525 (69)	140 (70)	0.77	799 (70)	31 (78)	0.38
Work status, n (%)			<0.001			0.23
Full- or part-time	1051 (48)	52 (26)		697 (61)	19 (48)	
Disabled	234 (11)	31 (16)		151 (13)	7 (18)	
Other	921 (42)	116 (58)		296 (26)	14 (35)	
Compensation, n (%)§	275 (12)	18 (9)	0.19	202 (18)	4 (10)	0.30
Body mass index; mean (SD)¶	28.3 (5.6)	32.9 (6.3)	<0.001	27.8 (5.4)	32.2 (7.3)	<0.001
Smoker, n (%)	375 (17)	20 (10)	0.015	275 (24)	7 (18)	0.45
Comorbidities, n (%)						
Hypertension	598 (27)	128 (64)	<0.001	143 (12)	20 (50)	<0.001
Stroke	25 (1)	11 (6)	<0.001	3 (0)	2 (5)	<0.001
Diabetes	25 (1)	199 (100)	<0.001	8 (1)	40 (100)	<0.001
Osteoporosis	132 (6)	15 (8)	0.47	16 (1)	2 (5)	0.24
Cancer	101 (5)	17 (9)	0.021	21 (2)	3 (8)	0.054
Depression	279 (13)	30 (15)	0.38	137 (12)	4 (10)	0.90
Anxiety	125 (6)	12 (6)	0.96	76 (7)	0 (0)	0.18
Drug dependency	10 (0)	0 (0)	0.71	6 (1)	0 (0)	0.50
Heart problem	289 (13)	59 (30)	<0.001	57 (5)	4 (10)	0.29
Lung problem	117 (5)	20 (10)	0.009	44 (4)	0 (0)	0.40
Stomach problem	364 (16)	50 (25)	0.003	135 (12)	7 (18)	0.40
Bowel or intestinal problem	185 (8)	24 (12)	0.10	75 (7)	5 (12)	0.25
Liver problem	28 (1)	3 (2)	0.97	13 (1)	0 (0)	0.92
Kidney problem	69 (3)	15 (8)	0.002	26 (2)	2 (5)	0.56
Blood vessel problem	79 (4)	13 (7)	0.059	15 (1)	1 (2)	0.96
Nervous system problem	46 (2)	7 (4)	0.29	18 (2)	0 (0)	0.89
Joint problem	801 (36)	110 (55)	<0.001	212 (19)	9 (22)	0.67

SpS			DS		
Not Diabetic, n = 538	Diabetic, n = 89	P†	Not Diabetic, n = 524	Diabetic, n = 70	P†
64.3 (12)	67.2 (9.3)	0.028	65.9 (10.4)	67.3 (9.1)	0.28
211 (39)	37 (42)	0.76	362 (69)	45 (64)	0.50
512 (95)	87 (98)	0.41	511 (98)	69 (99)	0.90
459 (85)	69 (78)	0.087	446 (85)	53 (76)	0.066
343 (64)	54 (61)	0.66	353 (67)	41 (59)	0.18
107 (20)	15 (17)	0.60	120 (23)	17 (24)	0.91
381 (71)	62 (70)	0.92	345 (66)	47 (67)	0.93
		0.009			0.043
179 (33)	18 (20)		175 (33)	15 (21)	
43 (8)	14 (16)		40 (8)	10 (14)	
316 (59)	57 (64)	309 (59)	45 (64)		
37 (7)	10 (11)	0.22	36 (7)	4 (6)	0.91
29 (5.3)	32.9 (6.3)	<0.001	28.7 (6)	33.2 (5.9)	<0.001
53 (10)	9 (10)	0.91	47 (9)	4 (6)	0.49
233 (43)	55 (62)	0.002	222 (42)	53 (76)	<0.001
11 (2)	2 (2)	0.78	11 (2)	7 (10)	0.001
7 (1)	89 (100)	<0.001	10 (2)	70 (100)	<0.001
52 (10)	8 (9)	0.99	64 (12)	5 (7)	0.30
40 (7)	8 (9)	0.77	40 (8)	6 (9)	0.97
61 (11)	9 (10)	0.87	81 (15)	17 (24)	0.09
23 (4)	7 (8)	0.23	26 (5)	5 (7)	0.63
2 (0)	0 (0)	0.66	2 (0)	0 (0)	0.56
136 (25)	29 (33)	0.19	96 (18)	26 (37)	<0.001
39 (7)	9 (10)	0.47	34 (6)	11 (16)	0.012
115 (21)	24 (27)	0.30	114 (22)	19 (27)	0.39
71 (13)	15 (17)	0.45	39 (7)	4 (6)	0.78
9 (2)	1 (1)	0.94	6 (1)	2 (3)	0.54
26 (5)	3 (3)	0.74	17 (3)	10 (14)	<0.001
31 (6)	7 (8)	0.60	33 (6)	5 (7)	0.99
10 (2)	3 (3)	0.60	18 (3)	4 (6)	0.54
292 (54)	54 (61)	0.31	297 (57)	47 (67)	0.12

(continued)

TABLE 1. (Continued)

Characteristics	All Analyzed Patients (IDH + SpS + DS)			IDH		
	Not Diabetic, n = 2207	Diabetic, n = 199	P†	Not Diabetic, n = 1145	Diabetic, n = 40	P†
Other**	696 (32)	83 (42)	0.004	486 (42)	40 (100)	<0.001
SF-36 scores, mean (SD)††						
Bodily pain	30.2 (20)	32.4 (20.3)	0.13	27.1 (20.1)	29.6 (20.2)	0.43
Physical functioning	36.8 (24.5)	29.3 (20.5)	<0.001	37.9 (25.5)	33.9 (25.2)	0.33
Vitality	41 (21.1)	37.8 (22.2)	0.038	38.5 (20.1)	36.1 (21.2)	0.46
Physical component summary	30.3 (8.5)	27.1 (8.2)	<0.001	30.6 (8.4)	28.8 (9.1)	0.18
Mental component summary	47.4 (11.8)	48.7 (12.3)	0.13	45.1 (11.6)	45.6 (11)	0.79
Oswestry Disability Index; mean (SD)‡‡	45.6 (20.2)	46.2 (19.2)	0.66	49.4 (21.3)	50.7 (21.9)	0.70
Sciatica Frequency Index (0–24); mean (SD)§§	14.9 (5.6)	14.5 (5.9)	0.28	15.9 (5.4)	15.6 (5.3)	0.79
Sciatica Bothersome Index (0–24); mean (SD)§§	15.1 (5.5)	14.6 (5.9)	0.25	15.6 (5.3)	14.8 (5.2)	0.35
Back Pain Bothersomeness (0–6); mean (SD)¶¶	4 (1.8)	4.3 (1.8)	0.035	3.9 (1.9)	3.9 (2)	0.89
Very dissatisfied with symptoms, n (%)	1656 (75)	132 (66)	0.009	920 (80)	28 (70)	0.16
Patient self-assessed health trend, n (%)			0.014			0.57
Getting better	249 (11)	14 (7)		173 (15)	6 (15)	
Staying about the same	864 (39)	66 (33)		518 (45)	15 (38)	
Getting worse	1092 (49)	119 (60)		453 (40)	19 (48)	
Treatment preference at baseline, n (%)			0.21			0.80
Preference for nonsurgery	772 (35)	80 (40)		378 (33)	15 (38)	
Not sure	415 (19)	40 (20)		190 (17)	7 (18)	
Preference for surgery	1015 (46)	79 (40)		574 (50)	18 (45)	

*Patients who report being told by their doctor that they have diabetes and also report that they are currently receiving treatment for diabetes.

†P values are from χ^2 test for categorical variables and t test for continuous variables.

‡Race or ethnic group was self-assessed. Whites and blacks could be either Hispanic or non-Hispanic.

§This category includes patients who were receiving or had applications pending for workers compensation, social security compensation, or other compensation.

¶The body mass index is the weight in kilograms divided by the square of the height in meters.

**Other indicates problems related to stroke, diabetes, osteoporosis, cancer, fibromyalgia, chronic fatigue syndrome, posttraumatic stress disorder, alcohol, drug dependence, heart, lung, liver, kidney, blood vessel, nervous system, hypertension, migraine, anxiety, stomach or bowel for IDH cohort and stroke, cancer, lung, fibromyalgia, chronic fatigue syndrome, posttraumatic stress disorder, alcohol, drug dependency, liver, kidney, blood vessel, nervous system, migraine, anxiety for SPS or DS cohort.

††The SF-36 scores range from 0 to 100, with higher score indicating less-severe symptoms.

‡‡The Oswestry Disability Index ranges from 0 to 100, with lower scores indicating less-severe symptoms.

§§The Sciatica Bothersomeness index range from 0 to 24, with lower scores indicating less-severe symptoms.

¶¶The Low Back Pain Bothersomeness Scale ranges from 0 to 6, with lower scores indicating less-severe symptoms

DS indicates degenerative spondylolisthesis; IDH, intervertebral disk herniation; SF-36, 36-Item Short Form Health Survey; SpS, spinal stenosis.

SpS			DS		
Not Diabetic, n = 538	Diabetic, n = 89	P†	Not Diabetic, n = 524	Diabetic, n = 70	P†
186 (35)	34 (38)	0.59	200 (38)	34 (49)	0.12
33.3 (20)	35.2 (18.3)	0.40	33.7 (18.6)	30.4 (22.4)	0.17
35.7 (24)	30.3 (17.9)	0.043	35.6 (22.4)	25.3 (20.1)	<0.001
42.9 (21.8)	41.2 (21.7)	0.49	44.6 (21.8)	34.4 (23.2)	<0.001
30.1 (8.8)	27.2 (7.4)	0.003	30 (8.2)	26 (8.7)	<0.001
49.2 (11.9)	51.1 (11.6)	0.17	50.4 (11.2)	47.5 (13.4)	0.043
41.9 (18.8)	44.6 (16.6)	0.19	41 (17.4)	45.6 (20.5)	0.041
14 (5.7)	13.1 (6)	0.18	13.8 (5.5)	15.5 (5.7)	0.013
14.5 (5.7)	13.5 (6)	0.16	14.5 (5.5)	15.8 (6)	0.076
4.1 (1.8)	4.1 (1.8)	0.73	4.2 (1.8)	4.8 (1.7)	0.014
374 (70)	54 (61)	0.12	362 (69)	50 (71)	0.79
		0.80			0.69
41 (8)	5 (6)		35 (7)	3 (4)	
174 (32)	29 (33)		172 (33)	22 (31)	
323 (60)	55 (62)		316 (60)	45 (64)	
		0.37			0.99
190 (35)	38 (43)		204 (39)	27 (39)	
104 (19)	17 (19)		121 (23)	16 (23)	
243 (45)	34 (38)		198 (38)	27 (39)	

$P < 0.001$) and for the PF score (29.3 for patients with diabetes *vs.* 36.8 for nondiabetic patients, $P < 0.001$). The impact of low back pain on a patient's daily function, as measured by the ODI, was not different for the two groups ($P = 0.66$).

INTERVERTEBRAL DISC HERNIATION

In the 1185 patients with IDH, only 40 IDH patients (3.4%) were diabetic (Table 1). These patients were significantly older than the nondiabetic population (mean age, 50.4 years *vs.* 41.5

years). Patients with diabetes had a significantly higher BMI ($P < 0.001$) and a significantly higher incidence of hypertension and stroke ($P < 0.001$ for both). Fewer IDH patients with diabetes than nondiabetic patients were working, but this was not statistically significant (48% *vs.* 61%, $P = 0.23$). There were no significant differences in SF-36 or ODI scores between the patients with diabetes and patients without diabetes.

Operative treatments, complications, and events were reviewed for the IDH subgroup (Table 2). After surgery, there

TABLE 2. Operative Treatments, Complications, and Events

	IDH		
	Not Diabetic, n = 768*	Diabetic, n = 24*	P
Specific procedures, n (%)†			
Decompression only	NA	NA	NA
Noninstrumented fusion	NA	NA	NA
Instrumented fusion	NA	NA	NA
Multi-level fusion	NA	NA	NA
Discectomy/decompression level, n (%)‡			
L2–L3	14 (2)	0 (0)	0.91
L3–4	27 (4)	0 (0)	0.71
L4–L5	297 (39)	14 (58)	0.093
L5–S1	438 (58)	11 (46)	0.34
Levels decompressed, n (%)			
None	NA	NA	NA
1	NA	NA	NA
2	NA	NA	NA
3+	NA	NA	NA
Operation time, min; mean (SD)	76.4 (37.4)	88 (37.9)	0.14
Blood loss, mL; mean (SD)	63.8 (102.9)	90.1 (72.1)	0.22
Blood Replacement, n (%)			
Intraoperative replacement	6 (1)	0 (0)	0.45
Postoperative transfusion	0 (0)	0 (0)	
Length of hospital stay, d; mean (SD)	0.97 (1)	1.1 (0.8)	0.55
Intraoperative complications, n (%)§			
Dural tear/spinal fluid leak	23 (3)	1 (4)	0.78
Vascular injury	0 (0)	0 (0)	
Nerve root injury	2 (0)	0 (0)	0.07
Other	3 (0)	0 (0)	0.17
None	741 (96)	23 (96)	0.70
Postoperative complications/events, n (%)¶			
Never-root injury	1 (0)	0 (0)	0.006
Wound dehiscence	0 (0)	0 (0)	
Wound hematoma	4 (1)	0 (0)	0.27
Wound infection	17 (2)	1 (4)	0.95
Other	26 (3)	1 (4)	0.71
None	719 (94)	22 (92)	0.95
Postoperative mortality, n (%)			
Death within 6 wk of surgery	0 (0)	0 (0)	
Death within 3 mo of surgery	1 (0.1)††	0 (0)	

SpS			DS		
Not Diabetic, n = 351*	Diabetic, n = 53*	P	Not Diabetic, n = 345*	Diabetic, n = 40*	P
		0.73			0.33
301 (88)	47 (90)		21 (6)	1 (2)	
19 (6)	3 (6)		74 (22)	6 (15)	
23 (7)	2 (4)		243 (72)	33 (82)	
14 (4)	2 (4)	0.76	79 (23)	10 (25)	0.92
124 (36)	21 (40)	0.65	35 (10)	9 (23)	0.039
239 (69)	37 (71)	0.93	168 (50)	19 (49)	0.97
315 (92)	51 (98)	0.17	331 (97)	39 (100)	0.53
135 (39)	17 (33)	0.45	104 (31)	9 (23)	0.43
		0.97		0.42	
7 (2)	1 (2)		3 (1)	1 (2)	
80 (23)	12 (23)		141 (41)	16 (40)	
110 (31)	15 (28)		126 (37)	11 (28)	
154 (44)	25 (47)		75 (22)	12 (30)	
127.3 (64.5)	140.3 (73.2)	0.19	205.7 (83.6)	216.8 (83.4)	0.43
299.5 (396.4)	373.6 (436.9)	0.21	571.6 (461.8)	688.4 (527.1)	0.14
32 (9)	7 (13)	0.52	111 (32)	20 (51)	0.029
15 (4)	5 (9)	0.21	66 (19)	14 (36)	0.028
3.1 (2.2)	3.8 (3.6)	0.051	5.7 (20.1)	5.1 (2.2)	0.83
32 (9)	5 (9)	0.85	38 (11)	2 (5)	0.36
0 (0)	0 (0)		0 (0)	1 (2)	0.19
0 (0)	0 (0)		0 (0)	0 (0)	
3 (1)	0 (0)	0.86	6 (2)	3 (8)	0.084
314 (90)	48 (91)	0.91	302 (88)	35 (88)	0.81
0 (0)	0 (0)		1 (0)	0 (0)	0.19
0 (0)	0 (0)		1 (0)	0 (0)	0.19
2 (1)	2 (4)	0.15	0 (0)	1 (3)	0.19
6 (2)	3 (6)	0.20	11 (3)	0 (0)	0.53
15 (4)	7 (13)	0.021	32 (9)	4 (10)	0.91
310 (90)	39 (74)	0.002	242 (71)	22 (56)	0.092
1 (0.3)	0 (0)	0.31	1 (0.3)	0 (0)	0.21
1 (0.3)	0 (0)	0.31	1 (0.3)	1 (2.3)	0.53

(Continued)

TABLE 2. (Continued)

	IDH		
	Not Diabetic, n = 768*	Diabetic, n = 24*	P
Additional surgeries, n (%)‡‡			
1-yr rate	43 (6)	3 (12)	0.13
2-yr rate	58 (7)	4 (17)	0.08
3-yr rate	65 (8)	4 (17)	0.13
4-yr rate	76 (10)	4 (17)	0.23
Recurrent disc herniation, n (%)	45 (6)	4 (17)	
Recurrent stenosis/progressive listhesis, n (%)	NA	NA	NA
Pseudarthrosis/fusion exploration,	NA	NA	NA
Complication or other, n (%)	21 (3)	0	
New condition, n (%)	9 (1)	0	

*Surgical information was available for 768 IDH patients without diabetes and 24 IDH patients with diabetes, 351 SpS patients without diabetes and 53 SpS patients with diabetes, and 345 DS patients without diabetes and 40 DS patients with diabetes.

†Specific procedure data was available for 343 SpS patients without diabetes and 52 SpS patients with diabetes, and 338 DS patients without diabetes and 40 DS patients with diabetes.

#In IDH patients, discectomy level is recorded, and in DS and SpS patients, decompression level is recorded.

§No cases were reported of aspiration into the respiratory tract or operation at wrong level.

¶Complications or events occurring up to 8 wk after surgery are listed. There were no reported cases of bone-graft complication, cerebrospinal fluid leak, paralysis, cauda equina injury, and pseudarthrosis.

‡‡Patient died after heart surgery at another hospital, the death was judged unrelated to spine surgery.

##Rates of repeated surgery at 1, 2, 3, and 4 yr are Kaplan-Meier estimates. P values were calculated with the use of the log-rank test. Numbers and percentages are based on the first additional surgery if more than one additional surgery.

DS indicates degenerative spondylolisthesis; IDH, intervertebral disk herniation; SpS, spinal stenosis.

was one nerve-root injury among patients without diabetes and none among those with diabetes ($P = 0.006$). There was no difference in infection or wound dehiscence. There were no differences in operation time, blood loss or replacement, the length of stay, postoperative mortality, or additional surgeries for IDH patients.

In the IDH cohort, the nondiabetic group had significantly greater improvement ($P < 0.001$) with surgery when compared with nonoperative treatment in BP, PF, and ODI at 4 years (Figure 1). For all patients who had surgery, nondiabetic patients had significantly greater improvement than patients with diabetes in BP ($P < 0.04$) and PF ($P < 0.001$). Patients




Figure 1. Primary Outcomes Across 4 years for Diabetics and Non-Diabetics with Intervertebral Disc Herniation.

SpS			DS		
Not Diabetic, n = 351*	Diabetic, n = 53*	P	Not Diabetic, n = 345*	Diabetic, n = 40*	P
20 (6)	2 (4)	0.51	23 (7)	3 (7)	0.83
30 (8)	2 (4)	0.22	42 (12)	7 (17)	0.36
42 (12)	4 (7)	0.31	47 (13)	8 (20)	0.30
47 (13)	6 (11)	0.60	49 (14)	10 (24)	0.09
NA	N/A	NA	NA	NA	NA
19 (6)	4 (8)		16 (5)	3 (8)	
0 (0)	0 (0)		3 (0.9)	1	
17 (4.9)	1		22 (6.5)	3 (7.7)	
7 (2)	1		8 (2.4)	1	

with diabetes did not have significant improvement with surgery *versus* nonoperative treatment for BP, PF, or ODI. Outcomes for nonoperative treatment were not different between patients with diabetes and patients without diabetes for BP ($P = 0.78$), PF ($P = 0.64$), or ODI ($P = 0.48$).

At 4 years, among those who had surgery, there was a significantly higher proportion of nondiabetic patients who were working compared with the diabetic group ($P = 0.019$) (Table 3).

SPINAL STENOSIS

Of 627 patients with SpS (Table 1), 89 patients (14.2%) had diabetes. Patients with diabetes were significantly older than those without diabetes (mean age, 67 years *vs.* 64 years, $P = 0.028$). BMI was higher in patients with diabetes ($P < 0.001$). The patients with diabetes had a significantly higher incidence of hypertension ($P = 0.002$) (Table 1).

Initial ODI scores were not significantly different between the two groups, although physical functioning and physical component scores were significantly lower in patients with diabetes than in those without diabetes at baseline ($P = 0.043$ for PF and $P = 0.003$ for PCS). There was no difference in workers' compensation between the two groups, though significantly more nondiabetic patients worked than did patients with diabetes (33% *vs.* 20%, $P = 0.009$).

Nondiabetic patients encountered significantly fewer postoperative complications than patients with diabetes, 90% *versus* 74% ($P = 0.002$) (Table 2). There were more infections among the patients with diabetes (6%) than among the nondiabetic patients (2%), but the difference was not statistically significant ($P = 0.20$). Postoperative complications included wound infection and hematoma, headaches, nausea and vomiting, anemia, postoperative hypoxia and confusion, urinary retention, and prolonged drainage. There was no significant difference in operation time, blood loss, blood replacement, or additional surgeries, or postoperative mortalities.

Across 4 years, SpS patients without diabetes made significant gains, with surgery relative to nonoperative care in BP

($P < 0.001$), PF ($P < 0.001$), and ODI ($P < 0.001$) (Figure 2). There was no significant difference in improvement with surgery between patients without diabetes and patients with diabetes for BP ($P = 0.41$) or ODI ($P = 0.52$). Nondiabetic patients who underwent surgery improved marginally more in PF ($P = 0.062$) relative to patients with diabetes. The patients with diabetes also did make significant gains, with surgery relative to nonoperative care in BP ($P < 0.001$), PF ($P < 0.001$), and ODI ($P < 0.001$). Nondiabetic patients had significantly greater improvement with nonoperative care than patients with diabetes in PF ($P = 0.028$) and ODI ($P = 0.013$) but not with BP ($P = 0.17$). (Table 4)

DEGENERATIVE SPONDYLOLISTHESIS

There were 594 patients with DS, of which, 70 patients (11.8%) had diabetes (Table 1). Patients with diabetes were not significantly older than nondiabetic patients; (mean age, 67 years *vs.* 66 years, $P = 0.28$). BMI was higher in patients with diabetes ($P < 0.001$). Significant comorbidities, including hypertension ($P < 0.001$), stroke ($P = 0.001$), heart ($P < 0.001$), lung ($P = 0.012$), and kidney disease ($P < 0.001$), were significantly more common among patients with diabetes (Table 1).

Patients with diabetes worked less than nondiabetic patients ($P = 0.043$), though there was no significant difference in worker's compensation. Patients with diabetes had significantly lower function according to their SF-36 scores for PF ($P < 0.001$), vitality ($P < 0.001$), and PCS ($P < 0.001$). The ODI score ($P = 0.041$) was also significantly worse in patients with diabetes (Table 1).

Blood replacement was greater in patients with diabetes both intraoperatively and after surgery ($P = 0.029$ and $P = 0.028$). There were no significant differences in intraoperative or postoperative complications (Table 2).

Across 4 years, nondiabetic DS patients who underwent surgery experienced significantly greater improvement in ODI ($P < 0.001$), BP ($P < 0.001$), and PF ($P < 0.001$) than for nonsurgical treatments (Figure 3). Nondiabetic patients who had surgery made significantly greater gains than patients

TABLE 3. Subgroup Results from Adjusted* As-Treated Outcome Analysis by Diabetes for the Randomized and Observational Cohorts Combined Patients with Lumbar Intervertebral Disk Herniation

Outcome	Diabetes	1-Yr			2-Yr	
IDH (randomized & observational)		Surgical, Mean (SE)	Nonoperative, Mean (SE)	Treatment Effect,† (95% CI)	Surgical, Mean (SE)	Nonoperative, Mean (SE)
Primary outcomes						
SF-36 bodily pain (0–100), mean (SE)	Not diabetic	45.9 (0.9)	33.1 (1.2)	12.7 (9.9 to 15.5)	45.2 (0.9)	35.1 (1.2)
	Diabetic‡	41.4 (5.1)	34.5 (5.9)	6.8 (−7.7 to 21.3)	34.3 (5.1)	34.2 (6.2)
	<i>P</i> §	0.39	0.82	0.43	0.035	0.89
SF-36 physical function (0–100), mean (SE)	Not diabetic	44 (0.8)	28.5 (1.1)	15.5 (13 to 18.1)	43.5 (0.8)	30.5 (1.1)
	Diabetic	31 (4.7)	29.7 (5.4)	1.2 (−11.9 to 14.4)	26.6 (4.7)	29.2 (5.6)
	<i>P</i>	0.007	0.83	0.036	<0.001	0.83
Mental component summary (0–100), mean SE)	Not diabetic	7.6 (0.4)	4.4 (0.5)	3.2 (2.1 to 4.3)	6.4 (0.4)	4.4 (0.5)
	Diabetic	6.8 (2.1)	2 (2.4)	4.8 (−1 to 10.6)	6.1 (2.1)	6.4 (2.5)
	<i>P</i>	0.70	0.32	0.60	0.89	0.45
Oswestry Disability Index (0–100), mean (SE)	Not diabetic	−37 (0.7)	−23 (0.9)	−14 (−16.1 to −11.9)	−36.7 (0.7)	−24.8 (0.9)
	Diabetic	−32.3 (4)	−29 (4.5)	−3.3 (−14 to 7.5)	−27 (3.9)	−29.1 (4.7)
	<i>P</i>	0.25	0.19	0.054	0.015	0.37
Very/somewhat satisfied with symptoms (%)	Not diabetic	71.5	46.3	25.3 (19.2 to 31.3)	72.4	52.7
	Diabetic	70.2	59.8	10.4 (−19.7 to 40.5)	65.8	30.8
	<i>P</i>	0.89	0.28	0.37	0.53	0.18
Very/somewhat satisfied with care (%)	Not diabetic	92.4	83.7	8.8 (4.4 to 13.1)	90.7	80.1
	Diabetic	80.5	76	4.5 (−21.5 to 30.5)	85.3	66
	<i>P</i>	0.048	0.33	0.39	0.39	0.20
Self-rated progress, major improvement (%)	Not diabetic	80.2	56.3	23.9 (18.1 to 29.8)	75.7	60.8
	Diabetic	77.4	70.4	7 (−20.6 to 34.6)	63.2	65.8
	<i>P</i>	0.76	0.26	0.30	0.21	0.72
Work status, working (%)	Not diabetic	79.3	77.4	1.9 (−2.9 to 6.6)	78.7	78.9
	Diabetic	62.1	69.2	−7.1 (−34.6 to 20.3)	57.4	54.3
	<i>P</i>	0.044	0.30	0.38	0.016	0.015

*Adjusted for age, sex, body mass index, baseline score (for SF-36, ODI), and center.

†Treatment effect is the difference between the surgical and nonoperative mean change from baseline. Analysis is done by using a mixed model with a random subject intercept term. Treatment is a time-varying covariate where a patients' experience before surgery is attributed to the nonoperative arm, and time is measured from enrollment; his/her postsurgery outcomes are attributed to the surgical arm, and time is measured from the time of surgery.

‡Patients who report being told by their doctors that they have diabetes and also report that they are currently receiving treatment for diabetes.

§*P* values at each time-point are from multiple degree-of-freedom Wald tests.

with diabetes who had surgery with regard to BP ($P = 0.003$) and PF ($P = 0.002$). However, diabetic patients with DS who underwent surgery had significantly better results than those treated nonoperatively for BP ($P < 0.001$), PF ($P < 0.002$), and the ODI ($P < 0.001$). In contrast to surgical outcomes, the outcomes of nonoperative treatment were not signifi-

cantly different between patients with diabetes and patients without diabetes. (Table 5)

DISCUSSION

An understanding of the baseline differences between diabetic patients and nondiabetic patients in the SPORT trial may help

	3-Yr			4-Yr		
Treatment Effect, [†] (95% CI)	Surgical, Mean (SE)	Nonoperative, Mean (SE)	Treatment Effect, [†] (95% CI)	Surgical, Mean (SE)	Nonoperative, Mean (SE)	Treatment Effect, [†] (95% CI)
10.1 (7.2 to 13)	45.9 (1)	35 (1.3)	10.9 (7.8 to 13.9)	47.2 (1)	33.9 (1.4)	13.3 (10.2 to 16.5)
0.1 (−14.6 to 14.8)	33 (5.9)	25.9 (6.4)	7.1 (−9.2 to 23.4)	46.8 (5.6)	37.7 (6.9)	9.1 (−7.6 to 25.7)
0.19	0.031	0.16	0.66	0.95	0.58	0.62
13 (10.4 to 15.6)	43.5 (0.9)	30.8 (1.2)	12.8 (10.1 to 15.5)	44.4 (0.9)	30.3 (1.2)	14.1 (11.3 to 17)
−2.7 (−16 to 10.7)	35.2 (5.4)	27.1 (5.9)	8.1 (−6.6 to 22.8)	28.3 (5.2)	20.8 (6.2)	7.4 (−7.6 to 22.5)
0.024	0.13	0.54	0.54	0.002	0.13	0.39
1.9 (0.8 to 3.1)	6.3 (0.4)	4.1 (0.5)	2.2 (1 to 3.4)	6 (0.4)	4.5 (0.5)	1.5 (0.2 to 2.8)
−0.3 (−6.2 to 5.7)	3.9 (2.4)	5.1 (2.6)	−1.3 (−7.8 to 5.3)	6.2 (2.3)	2.2 (2.9)	4 (−3 to 10.9)
0.47	0.31	0.70	0.31	0.94	0.42	0.49
−11.9 (−14 to −9.7)	−36.8 (0.7)	−25.3 (1)	−11.5 (−13.8 to −9.3)	−37.4 (0.8)	−25.1 (1)	−12.3 (−14.7 to −10)
2.1 (−8.8 to 13)	−32.8 (4.5)	−22.1 (4.9)	−10.6 (−22.6 to 1.3)	−34.3 (4.3)	−29.9 (5.1)	−4.4 (−16.6 to 7.8)
0.013	0.37	0.52	0.89	0.48	0.36	0.21
19.7 (13.5 to 25.8)	71.6	52.8	18.8 (12.3 to 25.4)	73.5	50	23.4 (16.6 to 30.3)
35 (5,65)	64.1	50.7	13.4 (−22.5 to 49.3)	85.1	61.6	23.5 (−8.5 to 55.5)
0.48	0.57	0.89	0.76	0.32	0.41	0.79
10.6 (5.8 to 15.4)	88.6	75.1	13.5 (8 to 19)	90.5	77.9	12.6 (7 to 18.3)
19.3 (−8.8 to 47.4)	93.7	81.6	12.1 (−11.9 to 36.1)	80.4	82.8	−2.3 (−30.6 to 26)
0.80	0.50	0.58	0.82	0.22	0.71	0.29
14.8 (8.8 to 20.9)	73.5	57	16.4 (9.9 to 22.9)	76.6	54.5	22 (15.2 to 28.8)
−2.6 (−33.5 to 28.3)	62.7	64.3	−1.6 (−36.2 to 33.1)	67.6	65.8	1.9 (−32.9 to 36.6)
0.29	0.36	0.60	0.31	0.45	0.44	0.29
−0.2 (−5 to 4.5)	76.4	74.4	2.1 (−3.3 to 7.4)	77.5	73.2	4.3 (−1.4 to 9.9)
3 (−27.3 to 33.4)	72.3	63.8	8.5 (−21.4 to 38.4)	57.8	70.2	−12.5 (−44.2 to 19.3)
0.81	0.64	0.30	0.64	0.019	0.72	0.094

to explain the larger treatment effects seen for nondiabetic patients across diagnostic groups, as well as the dichotomy in surgical outcomes between the diabetic patients with IDH and those with SpS and DS. In this study, patients with diabetes were significantly older than nondiabetic patients with IDH and SpS. The average age of nondiabetic patients was 53 years, while patients with diabetes averaged 64 years old. Of the subgroups, IDH had the largest difference in age (50.4 years, diabetic patients *vs.* 41.5 years, nondiabetic patients). Diabetic patients with SpS and DS were only slightly older (67 years) than the nondiabetic patients (SpS, 64 years and DS, 66 years).

Among patients with diabetes in the SPORT population, there were lower mean baseline PF and vitality scores. Pa-

tients with diabetes had lower baseline PCS scores and worked less than nondiabetic patients. Surprisingly, there was no difference in pain levels between patients with diabetes and patients without diabetes, suggesting that pain may not be the reason for the lower functional level of the diabetic patients. Alternatively, the difference may be secondary to the older age, obesity, and greater frequency of comorbid conditions among the patients with diabetes.³³

Simpson *et al*¹¹ compared lumbar spine surgical outcomes between patients with diabetes and patients without diabetes. They retrospectively reviewed the outcomes of 62 age- and sex-matched patients with and without diabetes and combined subpopulations of lumbar disc disease and SpS. The

TABLE 4. Subgroup Results from Adjusted* As-Treated Outcome Analysis by Diabetes for the Randomized and Observational Cohorts Combined Patients with Lumbar Spinal Stenosis

Outcome	Diabetes	1-Yr			2-Yr	
		Surgical, Mean (SE)	Nonoperative, Mean (SE)	Treatment Effect,† (95% CI)	Surgical, Mean (SE)	Nonoperative, Mean (SE)
Primary outcomes						
SF-36 bodily pain (0–100), mean (SE)	Not diabetic	29.4 (1.4)	14 (1.6)	15.4 (11.6 to 19.3)	28.6 (1.4)	15.6 (1.7)
	Diabetic‡	34 (3.4)	15.6 (3.6)	18.4 (9.6 to 27.1)	27.6 (3.4)	5 (3.9)
	<i>P</i> §	0.21	0.68	0.54	0.77	0.013
SF-36 physical function (0–100), mean (SE)	Not diabetic	26 (1.3)	11.2 (1.5)	14.7 (11.1 to 18.3)	23 (1.3)	14.2 (1.6)
	Diabetic	23.1 (3.3)	9.3 (3.5)	13.8 (5.7 to 22)	16.9 (3.3)	1.7 (3.7)
	<i>P</i>	0.42	0.60	0.84	0.077	0.002
Mental component summary (0–100), mean (SE)	Not diabetic	3.9 (0.6)	2.4 (0.7)	1.5 (−0.1 to 3.1)	3.4 (0.6)	1.8 (0.7)
	Diabetic	1.7 (1.4)	2.1 (1.5)	−0.4 (−4.1 to 3.3)	4.5 (1.4)	−2.1 (1.6)
	<i>P</i>	0.14	0.83	0.36	0.45	0.026
Oswestry Disability Index (0–100), mean (SE)	Not diabetic	−21 (1)	−9.1 (1.2)	−11.9 (−14.7 to −9)	−20.6 (1)	−10.1 (1.2)
	Diabetic	−21.7 (2.6)	−6.3 (2.8)	−15.4 (−22 to −8.8)	−17.9 (2.6)	−2.5 (3)
	<i>P</i>	0.80	0.35	0.33	0.33	0.018
Very/somewhat satisfied with symptoms (%)	Not diabetic	68.9	28.7	40.2 (31.6 to 48.7)	71.6	29.1
	Diabetic	69.8	19.5	50.3 (32.6 to 67.9)	54.9	19.1
	<i>P</i>	0.90	0.26	0.34	0.036	0.29
Very/somewhat satisfied with care (%)	Not diabetic	86.2	67.2	19 (10.8 to 27.2)	83.5	66.7
	Diabetic	85.9	76	9.9 (−6.6 to 26.4)	79.9	62.3
	<i>P</i>	0.95	0.31	0.47	0.55	0.69
Self-rated progress major improvement (%)	Not diabetic	68	25.6	42.5 (34.2 to 50.7)	66.5	27.1
	Diabetic	72.1	19.5	52.6 (35.5 to 69.7)	47	26.1
	<i>P</i>	0.59	0.42	0.34	0.031	0.92

*Adjusted for age, sex, body mass index, work status, income, smoking status, self-assessed health trend at baseline, treatment preference, baseline score (for SF-36, ODI), baseline stenosis bothersomeness, and center.

†Treatment effect is the difference between the surgical and nonoperative mean change from baseline. Analysis is done by using a mixed model with a random subject intercept term. Treatment is a time-varying covariate where a patients' experience before surgery is attributed to the nonoperative arm, and time is measured from enrollment; his(her) postsurgery outcomes are attributed to the surgical arm, and time is measured from the time of surgery.

‡Patients who report being told by their doctors that they have diabetes and also report that they are currently receiving treatment for diabetes.

§*P* values at each time-point are from multiple degree-of-freedom Wald tests.

CI indicates confidence interval; OBS, observational; OR, odds ratio; RCT, randomized controlled trial; SF-36, 36-Item Short Form Health Survey; SpS, spinal stenosis.

authors found poorer clinical outcomes, more infection, and longer hospitalizations among patients with diabetes. The poorer outcomes observed in patients with diabetes who underwent surgery in the study by Simpson *et al*¹¹ are consistent with the findings seen in the present study in the IDH population but not with the positive outcomes that were seen in the SpS population. The study of Simpson *et al*¹¹ differs from our study in several important respects. It was retrospective in nature and combined patients with different spinal disorders. The average age of the all patients in their study was

63 years, whereas in the SPORT population, the nondiabetic patients averaged 52.8 years and the diabetic patients averaged 63.9 years. The older age of the patients might account for the increased infection rate in patients in the study by Simpson *et al*,¹¹ which was not seen in the diabetic patients with IDH or SpS who underwent surgery in SPORT. The study by Simpson *et al*¹¹ also used a modified outcome measure for cervical discectomy proposed by Odom, while our study relied on the SF-36 and the ODI to assess changes in pain and functional outcome.

	3-Yr			4-Yr		
Treatment Effect,† (95% CI)	Surgical, Mean (SE)	Nonoperative, Mean (SE)	Treatment Effect,† (95% CI)	Surgical, Mean (SE)	Nonoperative, Mean (SE)	Treatment Effect,† (95% CI)
13 (9,17.1)	28.1 (1.4)	15 (1.8)	13.2 (8.9 to 17.4)	27.2 (1.5)	13.8 (1.9)	13.4 (8.8 to 18.1)
22.5 (13.3 to 31.8)	31.3 (3.5)	9.1 (4.3)	22.2 (12.1 to 32.3)	22.3 (3.8)	10.3 (4.6)	12 (1 to 23)
0.06	0.40	0.21	0.10	0.23	0.49	0.81
8.9 (5.1 to 12.7)	21.9 (1.3)	11.1 (1.7)	10.7 (6.8 to 14.7)	21.3 (1.5)	11.1 (1.8)	10.2 (5.8 to 14.5)
15.2 (6.6 to 23.8)	14 (3.4)	1.5 (4.1)	12.5 (3.1 to 21.9)	12.9 (3.6)	11.3 (4.3)	1.6 (-8.6 to 11.8)
0.18	0.032	0.029	0.73	0.03	0.96	0.12
1.5 (-0.2 to 3.2)	2.9 (0.6)	1.4 (0.7)	1.5 (-0.3 to 3.3)	2.4 (0.7)	1 (0.8)	1.4 (-0.6 to 3.4)
6.6 (2.6 to 10.5)	0.3 (1.5)	-2.5 (1.8)	2.8 (-1.5 to 7.1)	0.4 (1.6)	-4.4 (1.9)	4.8 (0.1 to 9.4)
0.019	0.10	0.043	0.58	0.24	0.01	0.19
-10.5 (-13.5 to -7.5)	-18.9 (1.1)	-9.8 (1.3)	-9.1 (-12.3 to -6)	-18.9 (1.2)	-10.2 (1.4)	-8.7 (-12.2 to -5.3)
-15.4 (-22.3 to -8.4)	-16.1 (2.7)	-1.2 (3.3)	-14.9 (-22.5 to -7.3)	-15.9 (2.9)	-2.6 (3.5)	-13.3 (-21.6 to -5)
0.20	0.32	0.015	0.16	0.34	0.044	0.31
42.5 (33.5 to 51.5)	65.3	35.5	29.8 (19.7 to 39.9)	65.1	31.9	33.3 (22.4 to 44.1)
35.8 (15.8 to 55.9)	71.7	34.3	37.4 (14.9 to 59.9)	48.5	27.5	21.1 (-4.1 to 46.2)
0.77	0.42	0.90	0.54	0.11	0.71	0.50
16.8 (7.9 to 25.7)	84.1	59.5	24.6 (14.8 to 34.4)	79.8	62.5	17.3 (6.3 to 28.3)
17.6 (-3.6 to 38.8)	84.1	71.1	13 (-8.2 to 34.2)	77.8	73.8	4 (-19.8 to 27.9)
0.93	1	0.34	0.47	0.80	0.41	0.43
39.4 (30.6 to 48.2)	61.7	29.1	32.6 (23.1 to 42.2)	54	23.3	30.7 (20.5 to 40.9)
20.9 (0 to 41.7)	60.9	21.1	39.8 (19.1 to 60.6)	44.1	12.7	31.3 (10.6 to 52.1)
0.24	0.93	0.43	0.55	0.34	0.29	0.68




Figure 2. Primary Outcomes Across 4 Years for Diabetics and Non-Diabetics with Spinal Stenosis.

TABLE 5. Subgroup Results from Adjusted* As-Treated Outcome Analysis by Diabetes for the Randomized and Observational Cohorts Combined Patients with Lumbar Lumbar Degenerative Spondylolisthesis

Outcome	Diabetes	1-Yr			2-Yr	
DS (randomized & observational)		Surgical, Mean (SE)	Nonoperative, Mean (SE)	Treatment Effect,† (95% CI)	Surgical, Mean (SE)	Nonoperative, Mean (SE)
Primary outcomes						
SF-36 bodily pain (0–100), mean (SE)	Not diabetic	33 (1.3)	14 (1.5)	19 (15.4 to 22.7)	33.3 (1.3)	13.2 (1.6)
	Diabetic‡	29.9 (3.9)	12.5 (4.2)	17.3 (6.7 to 28)	21.9 (3.6)	11.6 (4.5)
	<i>P</i> §	0.44	0.74	0.77	0.003	0.74
SF-36 physical function (0–100), mean (SE)	Not diabetic	28.5 (1.3)	10.9 (1.4)	17.6 (14.2 to 21.1)	27.3 (1.3)	9.8 (1.6)
	Diabetic	27.8 (3.8)	8.6 (4.1)	19.2 (9.1 to 29.4)	16.1 (3.5)	5.5 (4.3)
	<i>P</i>	0.86	0.59	0.77	0.003	0.34
Mental component summary (0–100), mean (SE)	Not diabetic	2.8 (0.5)	1.5 (0.6)	1.3 (−0.2 to 2.9)	2.9 (0.5)	1.2 (0.7)
	Diabetic	4.9 (1.6)	1.5 (1.7)	3.4 (−1.1 to 7.8)	0.7 (1.5)	−1.8 (1.9)
	<i>P</i>	0.20	0.96	0.39	0.15	0.12
Oswestry Disability Index (0–100), mean (SE)	Not diabetic	−24.8 (1)	−7.9 (1.1)	−16.9 (−19.5 to −14.2)	−24.7 (1)	−7.7 (1.2)
	Diabetic	−27.5 (3)	−7.8 (3.2)	−19.7 (−27.6 to −11.9)	−18.5 (2.8)	−9.3 (3.4)
	<i>P</i>	0.39	0.96	0.50	0.033	0.66
Very/somewhat satisfied with symptoms (%)	Not diabetic	71.9	26.8	45 (37.3 to 52.8)	70.6	32.4
	Diabetic	81.4	30.8	50.6 (29.3 to 72)	61.4	30.4
	<i>P</i>	0.28	0.67	0.59	0.29	0.86
Very/somewhat satisfied with care (%)	Not diabetic	89.8	67.6	22.2 (14.7 to 29.7)	88.4	68.3
	Diabetic	90.3	75.2	15.1 (−4.2 to 34.4)	93.3	57.8
	<i>P</i>	0.92	0.44	0.67	0.36	0.36
Self-rated progress major improvement (%)	Not diabetic	74.8	24.7	50.1 (42.6 to 57.6)	74.7	23.6
	Diabetic	84.1	26.5	57.5 (37.2 to 77.9)	63.2	19.5
	<i>P</i>	0.23	0.86	0.50	0.18	0.68

*Adjusted for age, sex, body mass index, work status, income, smoking status, self-assessed health trend at baseline, treatment preference, baseline score (for SF-36, ODI), baseline stenosis bothersomeness, and center.

†Treatment effect is the difference between the surgical and nonoperative mean change from baseline. Analysis is done by using a mixed model with a random subject intercept term. Treatment is a time-varying covariate where a patients' experience before surgery is attributed to the nonoperative arm, and time is measured from enrollment; his/her postsurgery outcomes are attributed to the surgical arm, and time is measured from the time of surgery.

‡Patients who report being told by their doctors that they have diabetes and also report that they are currently receiving treatment for diabetes.

§*P* values at each time-point are from multiple degree-of-freedom Wald tests.

CI indicates confidence interval; OBS, observational; OR, odds ratio; RCT, randomized controlled trial; SF-36, 36-Item Short Form Health Survey; SpS, spinal stenosis.

Patients with diabetes are predisposed to a less-optimal outcome with surgery than are nondiabetic patients. Misdiagnosis may be an issue. Diabetic polyneuropathy and predisposition to peripheral nerve lesions may cloud the clinical picture. Vascular insufficiency is more common among patients with diabetes and may cause radiating pain with ambulation.³⁴ Both coexistent vascular compromise and secondary peripheral neurologic pathology may also affect the ability of the nerve roots to recover from surgical decompression.^{35,36}

Peripheral neuropathy and endurance deficits are common in persons with diabetes. The previous factors can af-

fect strength and proprioception which predispose patients to greater risk of falls and slower walking speed.^{37–39} This results in lower scores on outcome measures related to PF but not necessarily for BP.

Cinotti *et al*¹⁰ performed a retrospective study that looked at 25 patients with and without diabetes who underwent surgery for SpS. In this population, the outcome was successful in both groups, without any significant differences. However, the nondiabetic patients in the study were older than those with diabetes (71 years *vs.* 68 years), and the nondiabetic patients were selected for comparison because they had a higher

	3-Yr			4-Yr		
Treatment Effect, [†] (95% CI)	Surgical, Mean (SE)	Nonoperative, Mean (SE)	Treatment Effect, [†] (95% CI)	Surgical, Mean (SE)	Nonoperative, Mean (SE)	Treatment Effect, [†] (95% CI)
20.1 (16.2 to 24.1)	33.9 (1.4)	15.3 (1.8)	18.6 (14.5 to 22.8)	32.2 (1.5)	15.5 (1.9)	16.7 (12.2 to 21.2)
10.3 (−0.3 to 20.9)	22.7 (3.7)	13.5 (4.9)	9.2 (−2.3 to 20.7)	27.3 (4.2)	16.7 (5.3)	10.6 (−2.1 to 23.3)
0.085	0.004	0.74	0.13	0.27	0.83	0.37
17.4 (13.7 to 21.2)	26.3 (1.3)	8.7 (1.7)	17.6 (13.7 to 21.6)	27.7 (1.4)	7.8 (1.8)	19.9 (15.6 to 24.2)
10.6 (0.5 to 20.7)	14.8 (3.6)	9.9 (4.8)	4.9 (−6.3 to 16)	18.9 (4)	8.7 (5.2)	10.2 (−2.1 to 22.5)
0.21	0.002	0.81	0.032	0.037	0.87	0.14
1.6 (0 to 3.3)	2.7 (0.6)	0.5 (0.7)	2.2 (0.4 to 4)	2.7 (0.6)	0.4 (0.8)	2.4 (0.4 to 4.3)
2.5 (−2 to 7)	0.6 (1.5)	−2.1 (2.1)	2.7 (−2.2 to 7.6)	0.1 (1.8)	−2.9 (2.4)	3 (−2.7 to 8.7)
0.72	0.19	0.24	0.86	0.17	0.19	0.83
−17 (−19.9 to −14.1)	−22.6 (1)	−8.8 (1.3)	−13.8 (−16.9 to −10.8)	−23.5 (1.1)	−8 (1.4)	−15.6 (−18.9 to −12.3)
−9.2 (−17.2 to −1.3)	−16.9 (2.9)	−11.4 (3.6)	−5.5 (−13.9 to 3)	−20.3 (3.1)	−12.7 (4)	−7.6 (−17 to 1.9)
0.068	0.056	0.49	0.065	0.33	0.26	0.11
38.3 (29.6 to 47)	66.9	37	30 (20.3 to 39.6)	64.2	28.9	35.3 (25.3 to 45.4)
31 (7 to 55)	62.4	28.9	33.5 (7.7 to 59.2)	68.6	39.7	28.9 (−1 to 58.9)
0.60	0.63	0.51	0.81	0.67	0.44	0.71
20.1 (11.7 to 28.4)	88	65.4	22.6 (13.4 to 31.8)	86.1	66.9	19.2 (9 to 29.4)
35.5 (13.7 to 57.4)	86.7	86.9	−0.2 (−18 to 17.6)	91.7	63.6	28.1 (1.8 to 54.5)
0.19	0.83	0.024	0.08	0.38	0.80	0.41
51.1 (43.1 to 59.1)	72.1	23.8	48.3 (39.8 to 56.8)	66.9	19.1	47.8 (38.7 to 56.9)
43.8 (22.3 to 65.3)	65.5	30.7	34.8 (9 to 60.5)	76	32.5	43.5 (15.3 to 71.7)
0.68	0.44	0.55	0.35	0.34	0.28	0.75




Figure 3. Primary Outcomes Across 4 Years for Diabetics and Non-Diabetics with Degenerative Spondylolisthesis

rate of comorbidity to match the two groups. The SPORT patients with diabetes had a higher rate of comorbidity than those without diabetes. This might account for the smaller treatment effect with surgery for patients with diabetes compared with nondiabetic patients in SPORT.

Arinzon *et al*⁸ retrospectively reviewed 257 consecutive patients and found that surgical decompression for SpS improved pain levels and basic activities of daily living in patients with diabetes, though the results were better in nondiabetic patients. The patients with diabetes were compared with an age-matched nondiabetic group that was older than the diabetic group (72 years *vs.* 70 years). As in SPORT, there were higher rates of postoperative complications in the diabetic population but, unlike SPORT, patient satisfaction was less than that of the nondiabetic control group. Another study by Airaksinen *et al*⁷ found that diabetes was associated with a lower ODI score after surgery for SpS, which we did not see in SPORT patients with diabetes. Our study might have different outcomes because it was prospective and used different outcome measures.

SPORT patients with spondylolisthesis showed no difference in postoperative infection rate or nonunion between the patients with diabetes and patients without diabetes. Bendo *et al*⁹ retrospectively also found that clinical results with posterior arthrodesis, as well as complications, were similar between patients with diabetes and patients without diabetes. This population included patients with IDH and SpS. There was no difference in postoperative complication rates. Glassman *et al*¹⁴ retrospectively looked at an age-matched population and found that insulin-dependent and non-insulin-dependent patients with diabetes had increased complications, including infections, postoperative root lesions, and blood loss. There was also an increased rate of nonunion among patients with diabetes. Browne *et al*¹² retrospectively reviewed national inpatient data from 197,000 patients who had lumbar fusions. Diabetes was found to be associated with increased risk for postoperative complications, including nonroutine discharge, increased hospital charges, and the length of stay.

Higher rates of obesity, older age, and the higher incidence of other concurrent medical problems found in patients with diabetes may predispose patients to complications and prolonged hospitalization after surgery. Fang *et al*¹³ found that preoperative risk factors for infection included smoking, age greater than 60 years, diabetes, previous surgical infection, increased BMI, and alcohol abuse. Deyo *et al*⁴⁰ found that age was a significant factor in morbidity and mortality in lumbar spine surgery. Katz *et al*⁴¹ also found that patients with diabetes had a high incidence of comorbidity after lumbar spine decompression. Further research is necessary to learn about the interaction and importance of these individual risk factors in predisposing patients to less-optimal outcomes and higher complication rates.

The greatest limitation of this study was the small number of diabetic patients in the IDH subgroup. While the number of patients with diabetes is a reasonable representation of patients with diabetes in the IDH age range, the population of patients with diabetes in the IDH population is small. There-

fore, final conclusions on whether or not patients with diabetes improve with discectomy for IDH should be made with caution, and a future study with greater numbers of diabetic patients must be undertaken to come to a more-definitive conclusion. In addition, we do not have any information about the baseline or posttreatment status of the diabetes in these patients with regard to glycemic control. The type, chronicity, and degree of control of diabetes mellitus have an impact on neurologic and vascular sequelae of the disease. This in turn, would be expected to influence the diagnosis, treatment, outcomes, and potential complications of the treatment of spinal disorders. A prospective study that takes these factors into account would help to clarify the patients who are most likely to have the best outcome and fewest complications with the various surgical and nonsurgical treatments. Furthermore, a routine screen for diabetes with a 2-hour postprandial blood glucose for all patients might well have diagnosed more patients with diabetes than was seen in our study. This might have changed our baseline as well as our outcome and complication data.¹⁻³

CONCLUSION

This is the first prospective study to compare surgical and nonsurgical outcomes between diabetic patients and nondiabetic patients. Diabetic patients who underwent surgery for IDH did not make significant improvements in pain and function at 4 years. Both diabetic and nondiabetic patients with SpS and DS benefited from surgery with regard to alleviating pain and improving function. However, nondiabetic patients with SpS or DS made greater functional gains with surgical intervention than did patients with diabetes. Diabetic DS patients did not have as much improvement in pain with surgery as the nondiabetic DS population. Nonoperative treatment for nondiabetic SpS patients also resulted in significant gains relative to diabetic SpS patients with regard to function but not pain.

➤ Key Points

- ❑ SPORT patients with diabetes are older, have higher BMIs, and have more comorbidities than nondiabetic patients.
- ❑ Diabetic patients with intervertebral disc herniation did not make significant gains in pain and function with surgical intervention relative to diabetic patients who underwent nonoperative treatment.
- ❑ Diabetic patients with spinal stenosis or degenerative spondylolisthesis experienced greater improvements in pain and function with surgical intervention when compared with nonoperative treatment.
- ❑ Postoperative complications, but not postoperative infections, were more prevalent in patients with diabetes than in nondiabetic patients with SpS.
- ❑ There was an increase in postoperative and intraoperative blood replacement in diabetic patients with DS.

Acknowledgments

The authors thank the National Institute of Arthritis and Musculoskeletal and Skin Diseases (U01-AR45444) and the Office of Research on Women's Health, the National Institutes of Health, and the National Institute of Occupational Safety and Health, the Centers for Disease Control and Prevention.

References

1. Bailes BK. Diabetes mellitus and its chronic complications. *AORN J* 2002;76:266–76, 78–82; quiz 83–6.
2. Bloomgarden ZT. Diabetes complications. *Diab Care* 2004;27:1506–14.
3. DUBY JJ, Campbell RK, Setter SM, et al. Diabetic neuropathy: an intensive review. *Am J Health Syst Pharm* 2004;61:160–73; quiz 75–6.
4. Feldman EL, Russell JW, Sullivan KA, et al. New insights into the pathogenesis of diabetic neuropathy. *Curr Opin Neurol* 1999;12:553–63.
5. Guthrie RA, Guthrie DW. Pathophysiology of diabetes mellitus. *Crit Care Nurs Q* 2004;27:113–25.
6. Sheetz MJ, King GL. Molecular understanding of hyperglycemia's adverse effects for diabetic complications. *JAMA* 2002;288:2579–88.
7. Engelgau MM, Geiss LS, Saaddine JB, et al. The evolving diabetes burden in the United States. *Ann Intern Med* 2004;140:945–50.
8. Airaksinen O, Herno A, Turunen V, et al. Surgical outcome of 438 patients treated surgically for lumbar spinal stenosis. *Spine* 1997;22:2278–82.
9. Arinzon Z, Adunsky A, Fidelman Z, et al. Outcomes of decompression surgery for lumbar spinal stenosis in elderly diabetic patients. *Eur Spine J* 2004;13:32–7.
10. Bendo JA, Spivak J, Moskvich R, et al. Instrumented posterior arthrodesis of the lumbar spine in patients with diabetes mellitus. *Am J Orthop* 2000;29:617–20.
11. Cinotti G, Postacchini F, Weinstein JN. Lumbar spinal stenosis and diabetes. Outcome of surgical decompression. *J Bone Joint Surg Br* 1994;76:215–9.
12. Simpson JM, Silveri CP, Balderston RA, et al. The results of operations on the lumbar spine in patients who have diabetes mellitus. *J Bone Joint Surg Am* 1993;75:1823–9.
13. Browne JA, Cook C, Pietrobon R, et al. Diabetes and early postoperative outcomes following lumbar fusion. *Spine* 2007;32:2214–9.
14. Fang A, Hu SS, Endres N, et al. Risk factors for infection after spinal surgery. *Spine* 2005;30:1460–5.
15. Glassman SD, Alegre G, Carreon L, et al. Perioperative complications of lumbar instrumentation and fusion in patients with diabetes mellitus. *Spine J* 2003;3:496–501.
16. Wimmer C, Gluch H, Franzreb M, et al. Predisposing factors for infection in spine surgery: a survey of 850 spinal procedures. *J Spinal Disord* 1998;11:124–8.
17. American Diabetes Association. National diabetes fact sheet. <http://www.diabetes.org/diabetes-statistics.jsp>. Accessed May 27, 2005.
18. Gregg EW, Cadwell BL, Cheng YJ, et al. Trends in the prevalence and ratio of diagnosed to undiagnosed diabetes according to obesity levels in the U.S. *Diab Care* 2004;27:2806–12.
19. Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical versus nonsurgical treatment for lumbar degenerative spondylolisthesis. *N Engl J Med* 2007;356:2257–70.
20. Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical vs nonoperative treatment for lumbar disk herniation: the Spine Patient Outcomes Research Trial (SPORT) observational cohort. *JAMA* 2006;296:2451–9.
21. Weinstein JN, Tosteson TD, Lurie JD, et al. Surgical versus nonsurgical therapy for lumbar spinal stenosis. *N Engl J Med* 2008;358:794–810.
22. Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical versus nonoperative treatment for lumbar disc herniation: four-year results for the Spine Patient Outcomes Research Trial (SPORT). *Spine* 2008;33:2789–800.
23. Birkmeyer NJ, Weinstein JN, Tosteson AN, et al. Design of the Spine Patient Outcomes Research Trial (SPORT). *Spine* 2002;27:1361–72.
24. McHorney CA, Ware JE, Jr, Rogers W, et al. The validity and relative precision of MOS short- and long-form health status scales and Dartmouth COOP charts. Results from the Medical Outcomes Study. *Med Care* 1992;30:MS253–65.
25. Ware JE, Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992;30:473–83.
26. Fairbank JC, Couper J, Davies JB, et al. The Oswestry Low Back Pain Disability questionnaire. *Physiotherapy* 1980;66:271–3.
27. Weinstein JN, Tosteson TD, Lurie JD, et al. Surgical vs nonoperative treatment for lumbar disk herniation: the Spine Patient Outcomes Research Trial (SPORT): a randomized trial. *JAMA* 2006;296:2441–50.
28. Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical compared with nonoperative treatment for lumbar degenerative spondylolisthesis. four-year results in the Spine Patient Outcomes Research Trial (SPORT) randomized and observational cohorts. *J Bone Joint Surg Am* 2009;91:1295–304.
29. Lurie JD, Facett SC, Hanscom B, et al. Lumbar discectomy outcomes vary by herniation level in the Spine Patient Outcomes Research Trial. *J Bone Joint Surg Am* 2008;90(9):1811–19.
30. Park DK, An HS, Lurie JD, et al. Does multilevel lumbar stenosis lead to poorer outcomes?: a subanalysis of the Spine Patient Outcomes Research Trial (SPORT) lumbar stenosis study. *Spine (Phila Pa 1976)* 2010;35(4):439–46.
31. Pearson A, Blood E, Lurie J, et al. Degenerative spondylolisthesis versus spinal stenosis: does a slip matter? Comparison of baseline characteristics and outcomes (SPORT). *Spine (Phila Pa 1976)* 2010;35(3):298–305.
32. Tosteson TD, Hanscom B, Blood EA, et al. POSTER: Statistical Methods for Cross-Over in the SPORT Lumbar Disc Herniation Trial. Hong Kong, PRC: International Society for Study of the Lumbar Spine; 2007.
33. Resnick HE, Vinik AI, Schwartz AV, et al. Independent effects of peripheral nerve dysfunction on lower-extremity physical function in old age: the Women's Health and Aging Study. *Diab Care* 2000;23:1642–7.
34. Kreines K, Johnson E, Albrink M, et al. The course of peripheral vascular disease in non-insulin-dependent diabetes. *Diab Care* 1985;8:235–43.
35. Blau J, Logue V. Intermittent claudication of the cauda equina. *Lancet* 1961;2:1081–6.
36. Dyck PJ, Karnes JL, O'Brien P, et al. The spatial distribution of fiber loss in diabetic polyneuropathy suggests ischemia. *Ann Neurol* 1986;19:440–9.
37. Cimbiz A, Cakir O. Evaluation of balance and physical fitness in diabetic neuropathic patients. *J Diab Complications* 2005;19:160–4.
38. Gutierrez EM, Helber MD, Dealva D, et al. Mild diabetic neuropathy affects ankle motor function. *Clin Biomech (Bristol, Avon)* 2001;16:522–8.
39. Ozdirenc M, Biberoglu S, Ozcan A. Evaluation of physical fitness in patients with Type 2 diabetes mellitus. *Diab Res Clin Pract* 2003;60:171–6.
40. Deyo RA, Cherkin DC, Loeser JD, et al. Morbidity and mortality in association with operations on the lumbar spine. The influence of age, diagnosis, and procedure. *J Bone Joint Surg Am* 1992;74:536–43.
41. Katz JN, Lipson SJ, Larson MG, et al. The outcome of decompressive laminectomy for degenerative lumbar stenosis. *J Bone Joint Surg Am* 1991;73:809–16.