

## ***Ashbya gossypii* \_IEM\_ prep\_protocol**

Using Tannic Acid (TA), Uranyl Acetate (UA) and p-Phenylenediamine (PPD) to retain ultrastructure and antigenicity.

1. Allow *Ashbya* to settle. 15 ml conical tubes will work quite well for steps 1-3.
2. Remove as much fluid as possible and add fix: 10-15x volume of 3%PF/1%GTA/1%TA in 0.1 M NaCacodylate pH 7.2-7.4 (NaCac) for 2 hours at room temperature (RT). Swirl sample tubes every 15 minutes during fixation step. NOTE: Reduce % of GTA and increase % of PF to get best labeling count; e.g. any combination from 4%PF/0%GTA to 3%PF/1%GTA.
3. Replace with fresh fix, leave for one hour at RT with swirling, and then overnight on rotator at 4°C.
4. Remove as much fluid as possible and rinse in PBS.
5. Re-suspend in 2mls of solution A (100mM KPO<sub>4</sub>, pH7.5, 1.2 M sorbitol) and add 200 microliters of 10mg/ml zymolyase. Incubate with gentle rotation at 37 C until see 75% of the hyphae are phase dark (approximately 5-35 minutes). Wash 1X in Sol. A, spinning 2 min. at 2000rpm in a microfuge.
6. Remove as much fluid as possible and rinse in fix: 3%PF/1%GTA/1%TA in 40mM K<sub>2</sub>HPO<sub>4</sub>-KH<sub>2</sub>PO<sub>4</sub> buffer (pH6.7) with 0.5mM MgCl<sub>2</sub> (KPO<sub>4</sub>).
7. Spin sample down (1500rpm) and then:
8. Remove as much fluid as and re-fix in 3%PF/1%GTA/1%TA in 0.1 M NaCac. Swirl sample tubes every 15 minutes during fix step. Replace with fresh fix and store at 4°C if necessary.
9. Gently centrifuge (1800rpm for 2mins) to get sample to bottom and rinse in 0.1M NaCac.
10. Centrifuge 2000RPM for 2mins and then quench aldehydes 2X , over one hour, with 50mM Glycine in 0.1M NaCac.
11. Rinse once more in 0.1M NaCac for 10 minutes. Spin to 2-3000RPM for 2mins to get sample to bottom of tube.
12. Rinse in dH<sub>2</sub>O 2x 20 min each to remove all traces of NaCac buffer.
13. 'En-bloc stain' with 1% Uranyl Acetate in dH<sub>2</sub>O for 1 hour RT.
14. Rinse in dH<sub>2</sub>O.

15. Dehydrate through Ethanol series: 30% w 1%PPD, 50% w 1%PPD, 70% w 1%PPD, 85% w 1%PPD -- 30 minutes each on rotator at RT. Add 1% p-phenylenediamine (PPD) to each ethanol solution within 20-30 minutes of use. Solution will be pink to light orange. If you make it up too early, the PPD will react with Ethanol and turn dark. Don't worry if PPD doesn't dissolve completely. It still works.
16. 85% Ethanol w 1%PPD on rotator at RT --- 4 rinses over 1 hr.
17. Re-suspend in a 2:1 mixture of 85% Ethanol w 1%PPD: LR White MEDIUM resin. Place on a rotator for 2 hr at RT
18. Re-suspend in 1:1 85% Ethanol w 1%PPD: LR White resin. Place on rotator for one hour and then store over night at 4-6°C (NO rotator). NOTE: solutions will turn dark (green-black; don't worry; it still works to retain proteins and ultrastructure. Just make as many changes as necessary in next step to get the resin solution clear again.
19. Make up fresh 85% Ethanol /1%PPD. Re-suspend in 1:2 85%EtOH w 1%PPD: LR White MEDIUM resin. Place on rotator 1 hr. at RT.
20. Re-suspend in 100% LR White MEDIUM resin. Do as many initial changes as needed to remove all the ETOH/PPD; i.e. when the dark solution turns clear. Then 3 changes over 6hours, using rotator at RT. Then store over night at 4-6°C (no rotator).
21. Warm to RT and 4 changes of 100% LR White MEDIUM resin; one hr each RT.
22. Transfer to gelatin capsules; leave for 4 hour at RT.
23. Polymerize at 50 °C for 24hr.
24. Remove from heat, remove sample from gelatin capsule. Store at 4 °C.
25. Semi thin sections (@0.5microns) Stain: Toluidine Blue.
26. For Immuno labeling sections; see separate protocol.
27. Thin Section: stain: UAaq for 5'. LC for 2'. IEM stain: UAaq for 4'; LC for 10" or UAaq for 4'; NO LC.

#### References:

1. Ellis, A (2010) p-Phenylenediamine: An Adjunct to and a substitute for Osmium Tetroxide. Microscopy Today Sept 2010 48-49
2. Brorson, S-H, etal (1999) Increased yield of immunogold labeling of

epoxy sections by adding *para*-phenylenediamine in the tissue processing. *Micron* 30 1999 561–566

3. Phend, K.D, etal (1995) An Osmium-free Method of Epon Embedment That Preserves both Ultrastructure and Antigenicity for Post-embedding Immunocytochemistry' *J Histochem Cytochem* 1995 V 43(3), 283-292.
4. Sawaguchi, A. etal (2000) A simple contrast enhancement by potassium permanganate oxidation for Lowicryl K4M ultrathin sections prepared by high pressure freezing/freeze substitution. *J Microsc*, Vol. 2000 V201, Pt 1, 77-83.
5. Berrymann, M.A. etal (1990) An Enhanced Method for Post-embedding Immunocytochemical Staining Which Preserves Cell Membranes' *JHistochem Cytochem* 1990 Vol. 38. No. 2, 159-170, 1990
6. Sakai, T. etal (2005) Addition of Phosphotungstic acid to ethanol improves ultrastructure and antigenicity of tissue embedded in LR white Resin. *Arch. Histol Cytol* 2005 68 (5) 333-347
7. Berrymann, M.A. etal (1992) Effects of Tannic Acid on Antigenicity and Membrane Contrast in Ultra structural Immunocytochemistry. *J Histochem Cytochem* 1992 Vol. 40 (6) 841-817